

High-resolution optical spectroscopy of DY Cen: diffuse interstellar bands in a proto-fullerene circumstellar environment?

D. A. García-Hernández^{1,2}, N. Kameswara Rao^{3,4} and David L. Lambert⁴

ABSTRACT

We search high-resolution and high-quality VLT/UVES optical spectra of the hot R Coronae Borealis (RCB) star DY Cen for electronic transitions of the C₆₀ molecule and diffuse interstellar bands (DIBs). We report the non-detection of the strongest C₆₀ electronic transitions (e.g., those at ~ 3760 , 3980 , and 4024 Å). Absence of C₆₀ absorption bands may support recent laboratory results, which show that the ~ 7.0 , 8.5 , 17.4 , and 18.8 μm emission features seen in DY Cen - and other similar objects with PAH-like dominated IR spectra - are attributable to proto-fullerenes or fullerene precursors rather than to C₆₀. DIBs towards DY Cen are normal for its reddening; the only exception is the DIB at 6284 Å (possibly also the 7223 Å DIB) that is found to be unusually strong. We also report the detection of a new broad (FWHM ~ 2 Å) and unidentified feature centered at ~ 4000 Å. We suggest that this new band may be related to the circumstellar proto-fullerenes seen at infrared wavelengths.

Subject headings: astrochemistry — circumstellar matter — ISM: molecules — dust, extinction — Stars: chemically peculiar — Stars: individual: (DY Cen)

1. Introduction

Fullerenes and fullerene-related molecules have attracted much attention since their discovery in the laboratory (Kroto et al. 1985) due to their potential applications in su-

¹Instituto de Astrofísica de Canarias, C/ Via Láctea s/n, 38200 La Laguna, Spain; agarcia@iac.es

²Departamento de Astrofísica, Universidad de La Laguna (ULL), E-38205 La Laguna, Spain

³543, 17th Main, IV Sector, HSR Layout, Bangalore 560102, (Indian Institute of Astrophysics, Bangalore 560034,) India; nkrao@iiap.res.in

⁴The W.J. McDonald Observatory, University of Texas, Austin, TX 78712-1083, USA; dll@astro.as.utexas.edu

perconducting materials, optical devices, and medicine. In the astrophysical context, these complex molecules have been proposed as explanations for unidentified astronomical features such as the intense UV absorption band at 217 nm (e.g., Cataldo & Iglesias-Groth 2009) and the enigmatic diffuse interstellar bands (DIBs) (Herbig 1995, 2000; Iglesias-Groth 2007). The remarkable stability of fullerenes against intense radiation (e.g., Cataldo, Strazzulla & Iglesias-Groth 2009) has encouraged the idea that fullerenes should be present in the interstellar medium. Indeed, the 9577 and 9632 Å DIBs observed in a few reddened stars lie near two electronic transitions of the C_{60} cation observed in rare gas matrices (Foing & Ehrenfreund 1994) but the presence of fullerenes in astrophysical environments has been uncertain until recently, when the mid-IR spectral signatures of the C_{60} and C_{70} fullerenes were detected in several Planetary Nebulae (Cami et al. 2010; García-Hernández et al. 2010, 2011c).

Interestingly, García-Hernández et al. (2010) showed that contrary to theoretical and experimental expectation, fullerenes are efficiently formed in normal H-rich circumstellar environments. The prevailing laboratory view was that the synthesis of fullerenes is much more efficient under hydrogen-poor conditions (Kroto et al. 1985; De Vries et al. 1993). Thus, circumstellar envelopes around R Coronae Borealis (RCB) stars (see e.g., Lambert & Rao 1994 for a review) have been considered as promising environments for the formation of fullerene molecules (Goeres & Sedlmayr 1992). This is because the RCBs’ hydrogen deficiency together with the He- and C-rich character of the gas resembles the laboratory conditions where fullerenes are produced. However, this suggestion is not supported by our recent *Spitzer* observations of a large sample of RCB stars, which show that fullerene formation is inefficient in the highly H-deficient environments characteristic of most RCBs (García-Hernández et al. 2011a,b).

Surprisingly, the only exception among RCBs is the least H-deficient RCB star DY Cen (possibly also the second least H-deficient RCB star V854 Cen) that shows mid-IR emission features at ~ 7.0 , 8.5, 17.4, and 18.8 μm coincident with the IR transitions of C_{60} . Contrary to fullerene-containing PNe, the latter C_{60} mid-IR features in DY Cen are seen in conjunction with very strong polycyclic aromatic hydrocarbon (PAH) features. This striking difference between the IR spectra of fullerene-containing PNe and DY Cen is explained by Duley & Hu (2012) who attribute the ~ 7.0 , 8.5, 17.4, and 18.8 μm features in DY Cen to proto-fullerenes (or fullerene precursors) rather than to the C_{60} molecule. Based on their laboratory spectroscopy of hydrogenated amorphous carbon (HAC) nanoparticles, Duley & Hu (2012) suggest that these four IR features may be due to proto-fullerenes in sources such as reflection nebulae, RCB stars, and proto-PNe, which also show the unidentified 16.4 μm feature and other very strong PAH-like features. In light of the now suspect infrared identification of neutral C_{60} , we sought to detect the molecule’s electronic transitions in

absorption from DY Cen’s optical spectrum, as an additional check to the presence of C_{60} and see whether Duley & Hu (2012)’s “proto-fullerenes” suggestion could be substantiated.

2. VLT/UVES optical spectroscopy

High-resolution ($R \sim 30,000$) and high-quality (signal-to-noise ratio $S/N \geq 200$) optical ($\sim 3300\text{--}9450$ Å) spectroscopic observations of DY Cen ($V=12.7$, $E(B-V)=0.50$, see Table 1) were obtained in the period February–March 2010 at the European Southern Observatory (ESO). The UVES spectrograph at the ESO VLT (Paranal, Chile) was used with the 1.2” slit and the standard setting DIC2 (390+760). We obtained 11 individual exposures of 1800 s each, giving a total exposure time of 5.5 hours. The S/N in the continuum in the summed DY Cen’s spectrum is ~ 200 at 4000 Å and higher than 250 at wavelengths longer than 6000 Å. The observed spectra - processed by the UVES data reduction pipeline (Ballester et al. 2000) - were corrected for heliocentric motion and the stellar continuum was fitted by using standard astronomical tasks in IRAF.

As a comparison star for DY Cen in our search for C_{60} and DIBs, we selected the very nearby B0.5 Ia bright supergiant HD 115842, which only differs in Galactic longitude and latitude by $0^\circ.88$ and $1^\circ.46$, respectively. This is because HD 115842 has the same reddening ($E(B-V)=0.5$) as DY Cen and it may be located at a comparable distance, sampling similar ISM conditions. A comparison between HD 115842 and DY Cen will probably enlighten the environmental changes associated with DY Cen. Thus, HD 115842 was also observed on the same dates as DY Cen with the same VLT/UVES set-up. Table 1 compares DY Cen with the comparison star HD 115842 in terms of Galactic coordinates, spectral type, magnitude, and reddening. Finally, a spectrum of BD -9° 4395, an extreme helium (EHe) star that shows a photospheric spectrum similar to that of DY Cen, was obtained by us with the 2.7m Harlan Smith telescope and the Tull coudé spectrograph (Tull et al. 1995) at a spectral resolving power of 60,000.

3. A search for neutral C_{60}

We have inspected the optical spectrum of DY Cen in order to search for the presence of optical absorptions of neutral C_{60} . The allowed transitions at wavelengths below 4100 Å are much more intense than those of the Herzberg-Teller induced forbidden transitions at longer wavelengths (Leach 1992). The strongest allowed electronic transitions of neutral gas phase C_{60} molecules, as predicted from laboratory experiments, are located at 3760 ± 5 ,

3980 $\pm$ 0.5, and 4024.0 $\pm$ 0.5 Å with widths of 8, 6, and 4 Å, respectively (Sassara et al. 2001). The oscillator strength (f) of the $\sim$ 4024 C₆₀ band is similar to that at 3980 Å (Leach 1992), while the 3760 Å band has a five times higher oscillator strength (Braga et al. 1991). DY Cen’s optical spectrum is dominated by strong He I, C II, Ne I, O II, and N II absorption and variable emission lines together with nebular forbidden emission lines from [O II], [N II], [S II], etc. The presence of these absorption and emission lines (FWHM $\sim$ 1 Å) complicates the detection of broad and weak absorption features.

We can find no evidence for the presence of C₆₀ in absorption (or emission) at the wavelengths of the expected electronic transitions:

The 3760 $\pm$ 5 Å band: This, the strongest of the three in our bandpass, falls amongst a series of O II lines. These lines give consistent O abundances suggesting that none is does blended with a C₆₀ feature (Figure 1, left panel), even though its central wavelength is uncertain by $\pm$ 5 Å (Sassara et al. 2001). Note that the apparent additional absorption at 3756 Å in DY Cen is a stellar blend (Figure 1, left panel).

The 3980 $\pm$ 0.5 Å band: This coincides with a stellar C II line from a multiplet represented by other lines. A secondary contributor may be the Al III 5d ²D - 8f ²F° multiplet (Figure 1, right panel). The C II lines are present also in BD +9°4395. This coincidence and the line’s ‘stellar’ width show that C₆₀ is not a significant contributor. An interesting feature of the right panel of Figure 1 is the appearance of an unidentified broad feature in DY Cen at 4000 Å (FWHM $\sim$ 2 Å; equivalent width of 91 mÅ). It is to be noted here that the 4000 Å feature is real. First, a strong argument in favour of the 4000 Å feature being real is that it does not appear in the spectra of the comparison star taken with the same set-up and at the same time. Second, the 4000 Å band is clearly seen when reducing the corresponding *Echelle* order alone. This is not a known DIB; it’s not present in HD 115842 where all DIBs have a similar strength to those in DY Cen (see below) and neither is it listed by Hobbs et al. (2008). Identification as C₆₀ would demand a wavelength error of either 20 Å for the 3980 Å transition or 24 Å for the 4024 Å transition but Sassara et al. give the wavelength uncertainty as just $\pm$ 0.5 Å for both transitions.

The C₆₀ band at 4024.0 $\pm$ 0.5: This is irretrievably blended with the He I 4026 Å line and adjacent lines.

We estimate that the one-sigma detection limits on the column density from the 3760 and 3980 Å C₆₀ bands in our DY Cen spectra (S/N $\sim$ 200) are 4.6×10^{12} and 1.5×10^{13} cm⁻², respectively¹. It is of interest to compare this column density limit of about 10¹³ cm⁻²

¹1-sigma detection limits for the EQWs in our spectra scale as $\sim 1.064 \times \text{FWHM} / (\text{S/N})$ (see e.g., Hobbs et al. 2008).

with the number of C_{60} molecules calculated from the infrared emission features attributed to these molecules. This exercise is attempted next.

The IR features measure the number of molecules in excited vibrational states but the electronic transitions sought here depend on the molecules in their ground vibrational state. Analysis of the strengths of the IR features with the estimated A-values (Cami et al. 2010; García-Hernández 2011a) shows that the excitation temperature is about 600 K. With this temperature, we estimate the number of molecules in the ground state: $N(C_{60}) = 1.0 \times 10^{46} d^2$ where d , the distance to DY Cen, is given in kpc.

The column density pertinent to our non-detection of C_{60} depends on how the line of sight to the star intersects the C_{60} -containing part of the circumstellar shell. Assume a spherical shell centered on the star with inner and outer radii of R_{out} and R_{in} , respectively, with the line of sight running directly from R_{out} to R_{in} . Assume a uniform density of C_{60} molecules throughout the shell, then the density of molecules is $n(C_{60})$ where

$$n(C_{60}) = \frac{7.0 \times 10^{-41}}{R_{out}^3 - R_{in}^3} N(C_{60}) \quad (1)$$

where R_{out} and R_{in} are given in AU.

Shell radii are estimated from the run of dust temperature with distance from the star. The equilibrium temperature of a gray dust grain $T_d(R)$ in an optically thin circumstellar environment is given by

$$T_d(R) = \left(\frac{R_*}{2R} \right)^{0.5} T_* \quad (2)$$

where R_* is the stellar radius, R is the radial distance from the stellar center, and T_* is the stellar blackbody temperature (Kwok 2007, p. 314, Equation (10.32)). If stellar radii and temperature are estimated from the stellar luminosity $L/L_\odot$, the distance R is provided by

$$R = \frac{1}{430} \left(\frac{L}{L_\odot} \right)^{0.5} \left(\frac{T_\odot}{T_d(R)} \right)^2 \quad (3)$$

for R in AU.

Adopting $L/L_\odot = 10^4$, we find $T_d = 600$ K at $R = 21$ AU and $T_d = 270$ K at 104 AU where the former temperature is the excitation temperature of the C_{60} molecules and the latter temperature is the blackbody temperature of the dust. If we adopt $R_{in} \simeq 10$ AU and

$R_{out} \simeq 100$ AU, we obtain $n(\text{C}_{60}) = 0.7d^2 \text{ cm}^{-3}$ or $n(\text{C}_{60}) = 34 \text{ cm}^{-3}$ for $d = 7$ kpc and 17 cm^{-3} for $d = 5$ kpc. Then, the estimates for the C_{60} column density along the path ($R_{out} - R_{in}$) are $4.6 \times 10^{16} \text{ cm}^{-2}$ for 7 kpc and $2.3 \times 10^{16} \text{ cm}^{-2}$ for 5 kpc.

Such estimates are uncomfortably greater than the observed upper limit of around 10^{13} cm^{-2} . One possible explanation of this 1000-fold discrepancy is that the line of sight to DY Cen does not intersect the C_{60} -containing regions of the circumstellar envelope. An alternative view raises doubts about the IR detection of the C_{60} molecule in DY Cen (possibly also in V854 Cen; García-Hernández et al. 2011a), supporting the recent claim that the ~ 7.0 , 8.5 , 17.4 , and $18.8 \mu\text{m}$ features seen in astronomical sources with PAH-like dominated IR spectra should be attributed to proto-fullerenes or fullerene precursors rather than to C_{60} (Duley & Hu 2012). The latter seems to be consistent with the characteristics of the DIBs towards DY Cen, which are studied below.

4. Diffuse bands from fullerene precursors?

As we have mentioned above, we detect in DY Cen a broad unidentified feature centered at $\sim 4000 \text{ \AA}$, which is seen in DY Cen only (Fig. 1). Note that no diffuse interstellar bands (DIBs) are known at this wavelength (see e.g., Hobbs et al. 2008). To our knowledge, no molecule is known to exhibit a strong electronic transition at $\sim 4000 \text{ \AA}$ and this is corroborated by inspecting the NIST Chemistry WebBook Database². We speculate that this $4000 \text{ \AA}$ band may be related to the proto-fullerenes or fullerene precursors seen in the IR and that could be very abundant in the circumstellar envelope of DY Cen. Thus, the $4000 \text{ \AA}$ feature may represent a new and very unusual diffuse band (DIB).

We used the exhaustive list of DIBs provided by Hobbs et al. (2008) in HD 204827 to search for them in the spectrum of DY Cen. DIBs detected in DY Cen (and the comparison star HD 115842) are given in Table 2, where we give the central wavelength (after applying the heliocentric correction), full width at half-maximum (FWHM) in $\text{\AA}$, equivalent width (EQW) in $\text{m\AA}$, central depth (A_c), and the S/N ratio in the neighbouring continuum. Figure 2 compares a selection of DIBs in DY Cen with those in HD 115842. Table 2 (also Figure 2) shows that the DIBs towards DY Cen have about the same or slightly less (up to 30 per cent) strength, the same FWHM, and a similar radial velocity as those towards HD 115842. Since both stars have roughly the same interstellar reddening $E(\text{B-V})$, the similar profiles and strengths for their DIBs is not surprising. In addition, for the well studied DIBs at 5780 , 5797 , 5850 , 6196 , 6379 , and $6614 \text{ \AA}$, the normalised equivalent widths $\text{EQW}/E(\text{B-V})$

²See <http://webbook.nist.gov/chemistry/>.

in DY Cen agree very well with those measured in field stars by Luna et al. (2008) and in the prototype star HD 183143 (Herbig 1995).

The exceptions to the above trends are the DIBs at 6284 Å and at 7223 Å. The 7223 Å DIB is stronger in DY Cen than in HD 115824 as seen from the divided spectrum, but we could not estimate its total absorption because of the interference from the superposed telluric lines. Figure 3 (left panel) shows the region of 6284 Å for both stars. The spectra have been cleaned of telluric lines. It is clear that this DIB towards DY Cen is stronger than towards HD 115842, suggesting that the carrier of the 6284 Å DIB (along with 7223 Å) is different from the rest of the classical DIBs. Indeed, the strength of this DIB seems to be poorly correlated with the interstellar reddening. For example, Luna et al. (2008) and Herbig (1995) obtained $EQW/E(B-V)$ of 0.90 and 1.50 for field reddened stars and HD 183143, respectively, while we measure a value of 1.10 towards DY Cen; Hobbs et al. (2008) give a value of 0.41 in HD 204827. Curiously, the post-AGB star IRAS 06530–0213, which shows the strongest unidentified 21 μm feature (Zhang, Kwok & Hrivnak 2010), also shows an unusually strong 6284 Å DIB (see Luna et al. 2008). The 21 μm feature may be attributed to HACs or their decomposition products such as fullerene precursors or intermediate products (see e.g., García-Hernández 2012). It is to be noted here that examination of atomic and molecular lines towards the sight lines of DY Cen and HD 115842 show that ionized lines are enhanced in strength in the sight line towards DY Cen relative to that of HD 115824, which might suggest that the carrier(s) of the 6284 Å and 7223 Å DIBs could even have some contribution from ionized species.

Finally, we would like to remark that the very broad 4430 Å DIB in DY Cen is almost identical to that in the comparison star HD 115842. This is shown in the righthand panel of Figure 3 where we compare the 4400–4450 Å spectral region for both stars. The 4430 Å DIB has been linked to fullerenes bigger than C_{60} (e.g., C_{80} , C_{240} , C_{320} , and C_{540}) and/or buckyonions such as $C_{60}@C_{240}$ and $C_{60}@C_{240}@C_{540}$ (Iglesias-Groth 2007). Our finding also would be consistent with fullerenes and fullerenes-containing molecules not being especially overabundant towards DY Cen.

5. Concluding Remarks

Duley & Hu (2012) show that the infrared features at ~ 7.0 , 8.5, 17.4, and 18.8 μm detected in astronomical sources with PAH-like dominated IR spectra - such as those of the RCB stars DY Cen and V854 Cen (García-Hernández et al. 2011a), the Reflection Nebulae NGC 7023 and NGC 2023 (Sellgren et al. 2010) or the proto-PN IRAS 01005+7910 (Zhang & Kwok 2011) should be attributed to fullerene precursors (or proto-fullerenes) rather than

to C_{60} . Our non detection of neutral C_{60} in DY Cen may support this claim . Thus, the carrier of the new 4000 Å absorption band detected in DY Cen may be intimately related with the fullerene precursors seen at infrared wavelengths. Duley & Hu (2012) suggest that fullerene precursors contain pentagonal carbon rings; the infrared emission features at ~ 7.0 and $16.4 \mu\text{m}$ usually detected in objects with PAH-like dominated IR spectra are characteristics of pentagonal rings (Moutou et al. 2000). Pentagonal carbon rings are present in carbon nanoparticles (e.g., HACs; Duley & Hu 2012) and nanotubes although these have less perfect structures than C_{60} and other fullerenes.

In summary, the non detection of neutral C_{60} in the high-quality VLT/UVES DY Cen’s spectrum may support recent experimental work, showing that the ~ 7.0 , 8.5 , 17.4 , and $18.8 \mu\text{m}$ IR features seen in sources with PAH-like dominated spectra have to attributed to proto-fullerenes rather than to neutral C_{60} . In addition, the new 4000 Å DIB reported here (possibly also the carriers of the classical 6284Å and 7223Å DIBs) may be related to fullerene precursors; an organic compound containing pentagonal rings. These pentagonal carbon rings are usually present in HAC nanoparticles and nanotubes, suggesting that they may be intimately related with the formation process of fullerenes.

This work is based on observations obtained with the ESO Very Large Telescope at Paranal under the Director’s Discretionary Time (DDT) programme 284.D- 5048(A) (PI: D. A. García-Hernández). D.A.G.H. acknowledges support provided by the Spanish Ministry of Economy and Competitiveness under grant AYA–2011–27754. D.A.G.H. also thanks his son Mateo for his great patience during the realization of this work. D.L.L. also wishes to thank the Robert A. Welch Foundation of Houston, Texas for support through grant F-634.

Facilities: VLT:UVES.

REFERENCES

- Ballester, P., Modigliani, A., Boitquin, O. et al. 2000, Messenger, 101, 31
- Braga, M., Larsson, S., Rosen, A., & Volosov, A. 1991, A&A, 245, 232
- Cami, J., Bernard-Salas, J., Peeters, E., & Malek, S. E. 2010, Science, 329, 1180
- Cataldo, F., & Iglesias-Groth, S. 2009, MNRAS, 400, 291
- Cataldo, F., Strazzulla, G., & Iglesias-Groth, S. 2009, MNRAS, 394, 615
- De Marco, O. et al. 2002, AJ, 123, 3387

- De Vries, M. S., Reihs, K., Wendt, H. R. et al. 1993, *Geochim. Cosmoc. Acta* 57, 933
- Duley, W. W., & Hu, A. 2012, *ApJ*, 745, L11
- Foing, B. H., & Ehrenfreund, P. 1994, *Nature*, 369, 296
- García-Hernández, D. A., Manchado, A., García-Lario, P. et al. 2010, *ApJ*, 724, L39
- García-Hernández, D. A., N. K., & Lambert, D. L. 2011a, *ApJ*, 729, 126
- García-Hernández, D. A., N. K., & Lambert, D. L. 2011b, *ApJ*, 739, 37
- García-Hernández, D. A., Iglesias-Groth, S, Acosta-Pulido, J. A. et al. 2011c, *ApJ*, 737, L30
- García-Hernández, D. A. 2012, in *“Planetary Nebulae: An Eye to the Future”*, Proc. of the IAU, IAU Symposium, 283, p. 148
- Goeres, A., & Sedlmayr, E. 1992, *A&A*, 265, 216
- Herbig, G. H. 1995, *ARA&A*, 33, 19
- Herbig, G. H. 2000, *ApJ*, 542, 334
- Hobbs, L. M., York, D. G., Snow, T. P. et al. 2008, *ApJ*, 680, 1256
- Hunter, I. et al. 2006, *MNRAS*, 367, 1478
- Iglesias-Groth, S. 2007, *ApJ*, 661, L167
- Krelowski, J., Beletsky, Y., Galazutdinov, G. A. et al. 2010, *ApJ*, 714, L64
- Kroto, H. W., Heath, J. R., O'Brien, S. C. et al. 1985, *Nature*, 318, 162
- Kwok, S. 2007, *Physics and Chemistry of the Interstellar Medium*, (Sausalito: Univ. Science Books)
- Lambert, D. L. & Rao, N. K. 1994, *JApA*, 15, 47
- Leach, S. 1992, *Chem. Phys.* 160, 451
- Luna, R., Cox, N. L. J., Satorre, M. A. et al. 2008, *A&A*, 480, 133
- Moutou, C., Verstraete, L., Léger, A. et al. 2000, *A&A*, 354, L17
- Sassara, A., Zerza, G., Chergui, M. & Leach, S. et al. 2001, *ApJ*, 135, 263
- Sellgren, K., Werner, M. W., Ingalls, J. G. et al. 2010, *ApJ*, 722, L54

Tull, R. G., MacQueen, P. J., Sneden, C., & Lambert, D. L. 1995, *PASP*, 107, 251

Zhang, Y., & Kwok, S. 2011. *ApJ*, 730, 126

Zhang, Y., Kwok, S., & Hrivnak, B. J. 2010, *ApJ*, 725, 990

Table 1: Observational parameters of DY Cen.

Star	l	b	V	B-V	E(B-V)	Sp. Type	M_V	Distance (pc)	Ref
DY Cen	307.958	8.293	12.7	0.33	0.47-0.50	B0 Ia		4800	1
HD115842	307.080	6.834	6.04	0.30	0.50	B0.5 Ia	-8.0	3581	
							-6.9	2157	2

1:- De Marco et al(2002), 2: Krelowski et al (2010), Hunter et al (2006)

Table 2: Diffuse Interstellar Bands in DY Cen and HD 115842^a

<u>DY Cen</u>					<u>HD 115842</u>				
λ_c (Å)	FWHM (Å)	EQW (mÅ)	Ac	S/N	λ_c (Å)	FWHM (Å)	EQW (mÅ)	Ac	S/N
4000.38	1.92	91	0.04	188					412
4429.66	21.10	1034	0.05	273	4428.50	23.00	932	0.04	436
5780.46	2.14	256	0.10	209	5780.39	2.10	257	0.11	307
5796.92	0.91	72	0.07	199	5796.90	0.89	105	0.11	264
5849.67	0.82	32	0.02	272	5849.58	0.83	43	0.05	330
6065.11	0.61	6	0.01	312	6065.00	0.64	7	0.01	635
6089.62	0.53	7	0.01	282	6089.59	0.53	13	0.02	490
6112.7	1.00	12	0.01	300	6112.97	0.62	12	0.02	490
					6116.62	0.94	8	0.01	490
6139.75	0.82	5	0.01	267	6139.72	0.59	8	0.01	570
6195.80	0.60	27	0.04	284	6195.80	0.50	33	0.06	636
6202.81	1.28	62	0.04	336	6202.83	1.35	64	0.04	636
6233.79	0.79	10	0.01	336	6233.80	0.66	14	0.02	560
6269.70	1.21	36	0.02	276	6269.80	1.38	70	0.04	582
6283.53	4.29	543	0.12	276	6283.75	4.43	413	0.09	582
6375.97	0.92	13	0.02	273	6375.91	0.90	29	0.03	340
6379.25 ^b	1.04	53			6379.04	0.64	70	0.10	340
6597.26	0.53	8	0.01	295	6597.19	0.58	9	0.01	378
6613.41	1.03	83	0.08	295	6613.45	1.04	149	0.14	304
6660.60	0.64	12	0.02	314	6660.53	0.80	39	0.05	265
7223.75									
8620.88	3.75	95	0.02	106	8620.94	3.92	105	0.02	243

^a The $3\text{-}\sigma$ errors in the EQWs scale like $\sim 3 \times \text{FWHM} / (\text{S/N})$ while we estimate that the FWHMs are precise to the 0.03 Å level; ^b blended with N II 6379.98 Å




Fig. 1.— The spectra of DY Cen (in red) and HD 115842 (in blue) around 3760 Å (left panel) and 4000 Å (right panel; where the eHe star BD -9° 4395 is also displayed in green) are shown. The expected positions of the C_{60} features are marked on top of the spectra.. The FWHM of the features are indicated by the dotted line. Note that there is no evidence (additional absorption) in DY Cen for the presence of the neutral C_{60} features at 3760, 3980, and 4024 Å. However, there is an additional absorption band at 4000 Å in DY Cen (marked by a dark line and written DIB4000), which is not present in either HD 115842 or in the extreme He star BD -9° 4395 that is expected to show similar spectrum as DY Cen.




Fig. 2.— Profiles of six DIBs of various strengths and widths are plotted with respect to the LSR radial velocity. Note that the profiles are shifted vertically for clarity. The [Fe II] profile at the bottom shows the DY Cen’s systemic radial velocity. Note that the strengths and shapes of 5780 and 6203 Å DIBs are almost identical for both stars. The other profiles show that DIBs towards the DY Cen sight of line are weaker than towards HD 115842. To convert the velocity scale to heliocentric, 4.6 and 5.1 km s^{-1} should be added to the LSR velocities of DY Cen and HD 115842, respectively.




Fig. 3.— Spectral regions around the DIBs at 6284 Å (left panel) and 4428 Å (right panel) in DY Cen (in red) and HD 115842 (in blue). The telluric line corrected spectrum of both stars are shown by smooth lines (red and blue). The not fully telluric corrected spectrum with emission of [O I] line in DY Cen is shown in green. It is obvious that the 6284 Å DIB is stronger in DY Cen spectrum than towards HD 115842 while the DIB at 4428 Å (right panel) is of similar strength in both stars.